

## ABSTRACTS (most recent first)

A99) Sputtek, A., Nowicki, B., Rowe A.W., Kühnl, P.  
Long-term cryopreservation of human peripheral blood progenitor cells (PBPC):  
Influence of storage temperature (-80 °C vs. <-170 °C) on cell recovery, membrane  
integrity, and clonogenicity  
Cryobiology 49, 314 (2004)

A98) Rowe, A.W., Sputtek, A.  
Cryopreservation of blood cells. A brief overview  
Cryobiology 49, 310-311 (2004)

A97) Rowe, A.W., Sputtek, A.  
Autologous transfusion of cryopreserved blood cells  
Low Temp. Med. 30, 100 (2004)

A96) Sputtek, A.  
Hydroxyethyl Starch (HES) – An alternative for the freezing of human red blood cells  
Low Temp. Med. 30, 99 (2004)

A95) Sputtek, A.  
Protective mechanisms of penetrating and non-penetrating cryoprotectants  
Low Temp. Med. 30, 80 (2004)

A94) Sputtek, A., Nowicki, B., Rowe A.W., Kühnl, P.  
-80 °C or below –170 °C: Which temperature do we need for long-term storage of  
human peripheral blood progenitor cells (PBPC)?  
Tranfus Med Hemother 31 (Suppl 3), 10 (2004)

A93) Nowicki, B., Sputtek, A., Kühnl, P.  
Influence of storage temperature (-80 °C vs. <-150 °C) on cell recovery, membrane  
integrity and clonogenicity after long-term storage of human PBPC  
Cytotherapy 6, 403 (2004)

A92) Nowicki, B., Sputtek, A., Kühnl, P.  
Effect of long-term storage at –80 °C versus below –150 °C of human peripheral  
blood progenitor cells on cell recovery, membrane integrity and hematopoietic  
potential  
Oncology 25 (Suppl 4), 112-113 (2002)

A91) Sputtek, A., Benndorf, C., Rowe, A.W., Kühnl, P.  
Hematopoietic potential of cryopreserved peripheral blood stem cells after five hour  
post-thaw storage at 0 °C and 20 °C  
Low Temp. Med. 28, 111 (2002)

A90) Sputtek, A., Benndorf, C., Kühnl, P.  
Post-thaw storage of peripheral blood stem cell concentrates leads to a decrease of  
the hematopoietic potential and an increase of the danger of clotting  
Infus. Ther. Transfus. Med. 29 (Suppl 1), 25 (2002)

A89) Sputtek, A., Benndorf, C., Rowe, A.W., Kühnl, P.

Why frozen peripheral blood progenitor cell (PBPC) concentrates should be thawed at the patient's bedside  
Transfusion 42 (Suppl), 46S (2002)

A88) Sputtek, A., Benndorf, C., Rowe, A.W., Kühnl, P.  
Post-thaw storage of frozen peripheral blood stem cells decreases the hematopoietic potential  
Cryobiology 45, 248-249 (2002)

A87) Schulte am Esch J., Hamann, D., Soltau, M., Zante, B., Jungblut, M., Sputtek, A., Nierhaus, A., Hillert, C., Fischer, L., Rogiers, X.  
Ablagerung humaner Antikörper und Komplementaktivierung auf Schweine-Hepatozyten in einem klinisch angewendeten bio-artifiziellen Leberunterstützungssystem  
Transplantationsmed. Organ Dtsch. Transplantationsges. 13 (Suppl), 209 (2001)

A86) Sputtek, A.  
Cryopreservation and quality control of peripheral blood progenitor cells  
Clin. Lab. 47, 565 (2001)

A85) Sputtek, A., Hellstern, P.  
Frozen erythrocytes for immunizing Rh-negative plasma donors for anti-D production  
Infus. Ther. Transfus. Med. 28 (Suppl 1), 64 (2001)

A84) Siemensen, M.K., Brockmann, M.A., Geidel, K., Sputtek, A., Kühnl, P., Gutensohn, K.  
Quality control of platelet concentrates after cryopreservation compared to horizontally stored platelets using resonance thrombography  
Infus. Ther. Transfus. Med. 28 (Suppl 1), 57 (2001)

A82) Siemensen, M.K., Brockmann, M.A., Geidel, K., Sputtek, A., Kühnl, P., Gutensohn, K.  
Flow cytometric analysis of CD62p expression on the surface of cryopreserved platelet products  
Infus. Ther. Transfus. Med. 28 (Suppl 1), 55 (2001)

A81) Giese, L.-H., Pieper, N., Sputtek, A., Kühnl, P.  
Influence of the pre-freeze washing medium on postthaw recovery and postthaw storage stability of red cells using an automated cytopheresis device for collection and washing and a hydroxyethyl starch method for freezing  
Infus. Ther. Transfus. Med. 28 (Suppl 1), 55 (2001)

A80) Sputtek, A., Pieper, N., Giese, L.-H. Kühnl, P.  
Effect of electrolyte medium on the post-thaw recovery and post-thaw storage stability of erythrocytes (RBC) frozen in the presence of hydroxyethyl starch (HES)  
Cryobiology 43, 355-356 (2001)

A79) Schulte am Esch, J., Hamann, D., Soltau, M., Zante, B., Jungbluth, M., Sputtek, A., Nierhaus, A., Hillert, Ch., Fischer, L., Rogiers, X.  
Transplantationsmed. Organ. Dtsch. Transplantationsges. 13 (Suppl), 209 (2001)

A78) Sputtek, A., Benndorf, C., Eiermann Th., Kühnl, P.  
Stability and viability of cryopreserved peripheral blood stem cells (PBSC) during postthaw storage  
Infusionsther. Transfusionsmed. 26 (Suppl 1), 56 (1999)

A77) Sputtek, A.  
Cryopreservation of peripheral blood progenitor cells  
Infusionsther. Transfusionsmed. 26 (Suppl 1), 9 (1999)

A76) Sputtek, A., Voll, M., Kühnl, P.  
Post-thaw storage stability of hydroxyethyl starch-protected frozen red cells  
Cryobiology 37, 431 (1998)

A75) Sputtek, A., Kühnl, P.  
Cryopreservation of peripheral blood progenitor cells (PBPC)  
Cryobiology 37, 429 (1998)

A74) Sputtek, A.  
Kryokonservierung von hämopoetischen Stammzellen, Erythrozyten, Thrombozyten und Lymphozyten - Pause vom Leben?  
Infusionsther. Transfusionsmed. 25 (Suppl 1), 47 (1998)

A73) Gutensohn, K., Carrero, I., Lüdemann, K., Sputtek, A., Krüger, W., Kühnl, P.  
Semi-automated flow cytometric analysis of CD34-positive Cells in peripheral blood stem cell concentrates  
Infusionsther. Transfusionsmed. 25 (Suppl 1), 20 (1998)

A72) Löliger, C., Seyedin, F., Carrero, I., Göpfert, C., Gutensohn, G., Sputtek, A., Rauhöft, C., Fiedler, W., Kröger, N., Kühnl, P.  
Automated collection of peripheral blood progenitor stem cells using the Auto-PBSC software  
Blood 90, 330b, (1997)

A71) Sputtek, A.  
The role of penetrating and non-penetrating cryoprotectants in transfusion medicine  
In: Cryoprotectants in Medical Practice (Utilisation des cryoprotecteurs en médecine),  
Ed.: Institute International du Froid (IIF), ISBN 2-903633932, p 12, Paris, Frankreich (1997)

A70) Jetter, S., Sputtek, A., Hummel, K., Kühnl, P.  
Numerical recovery and membrane integrity of frozen/thawed human peripheral blood progenitor cells (PBPC) are unsuitable assays for the optimization of the cryopreservation procedure  
Infusionsther. Transfusionsmed. 24, 388 (1997)

A69) Sputtek, A.  
Hydroxyethyl starch, red cells and cryopreservation: facts and fictions  
Infusionsther. Transfusionsmed. 24, 381 (1997)

A68) Horn, E.-P., Sputtek, A., Voll, M., Lüdemann, K., Rudolf, B., Standl, Th., Schulte am Esch, J., Kühnl, P.

Effects of hydroxyethyl starch cryopreserved red blood cells on haemodynamics and tissue oxygenation  
Infusionsther. Transfusionsmed. 24, 375 (1997)

A67) Meuser, D., Sputtek, A., Kühnl, P.  
A cryopreservation procedure for human red blood cells (RBC) applicable for routine clinical practice using hydroxyethyl starch and liquid nitrogen  
Infusionsther. Transfusionsmed. 24, 374-375 (1997)

A66) Gutensohn, K., Garritsen, H.S.P., Cassens, U., Carrero, I., Sputtek, A., Alisch, A., Kühnl, P.  
Counting of residual leukocytes in platelet concentrates by quantitative flow cytometric bead assay  
Infusionsther. Transfusionsmed. 24, 284 (1997)

A65) Feldmann, C., Gutensohn, K., Sputtek, A.  
Cryopreservation of peripheral blood progenitor cells (PBPC) - Is there an influence of cooling rate and composition of the cryoprotective solution on progenitor subsets?  
Infusionsther. Transfusionsmed. 24, 240 (1997)

A64) Sputtek, A., Hummel, K., Eiermann, Th., Kühnl, P.  
Recovery of peripheral blood progenitor cells (PBPC) frozen in reference samples compared to the recovery in the simultaneously frozen bags for therapeutical use  
Infusionsther. Transfusionsmed. 24, 240 (1997)

A63) Sputtek, A., Gutensohn, K., Kühnl, P.  
Mechanismen der Kryoschädigung und -protektion bei isolierten Zellen  
Eur. J. Cell Biol. 74 (Suppl 45), 134 (1997); Optik 106 (Suppl 7), 134 (1997)

A62) Sputtek, A.  
Kyokonservierung von Erythrozyten mit Hydroxyethylstärke - Eine Alternative zur Flüssiglagerung in additiven Lösungen?  
Anästh. Intensivmed. 32 (Suppl 1), S29 (1997)

A61) Sputtek, A., Hammes, R., Warnken, U.H., Kühnl, P.  
A novel HPLC method for the determination of HES concentrations in plasma and urine after administration of HES protected frozen erythrocytes  
Cryo Letters 18, 72 (1997)

A60) Horn, E.-P., Sputtek, A., Rudolf, B., Standl, Th., Schulte am Esch, J.  
Cryopreservation of red blood cells using hydroxyethyl starch: a study on safety, tolerance, and oxygenation in patients  
Anesthesiology 85 (Suppl), A348 (1996)

A59) Gutensohn, K., Hummel, K., Krüger, W., Sputtek, A., Kühnl, P.  
Progressive alteration of platelet antigens during peripheral blood progenitor cell (PBPC) apheresis  
Ann. Hematol. 73 (Suppl 2), 160 (1996)

A58) Sputtek, A., Roos, D., Lubitz, B., Gutensohn, K., Müller, M., Kühnl, P.  
Einfluß des Bestrahlungsvorgangs auf Thrombozyten-Konzentrate

Infusionsther. Transfusionsmed. 23 (Suppl 3), 67 (1996)

A57) Löliger, C., Hummel, K., Gutensohn, K., Sputtek, A., Hassan, H. T., Kühnl, P.  
Biologische Eigenschaften von peripheren Blut-Progenitor-Zell-Präparaten für die  
allogene Transplantation

Infusionsther. Transfusionsmed. 23 (Suppl 3), 61 (1996)

A56) Löliger, C., Hummel, K., Gutensohn, K., Sputtek, A., Hiller, J., Kühnl, P.  
Thrombozytopenie bei einem gesunden Spender nach Apherese peripherer Blut-  
Progenitor Zellen für die allogene Transplantation

Infusionsther. Transfusionsmed. 23 (Suppl 3), 61 (1996)

A55) Hummel, K., Sputtek, A., Gutensohn, K., Löliger, C., Hassan, H. T., Zander. A.  
R., Kühnl, P.

Prädikativer Wert der Leukozytenkonzentration am Apheresetag für den Gehalt an  
CD34+-Zellen im PBSZ-Präparat bei Spendern

Infusionsther. Transfusionsmed. 23 (Suppl 3), 55 (1996)

A54) Sputtek, A., Jetter, S., Hummel, K., Kühnl, P.

Tiefgefrieren von peripheren Blutstammzellen - Welche Merkmale müssen geeignete  
Kryokonservierungsverfahren aufweisen?

Infusionsther. Transfusionsmed. 23 (Suppl 3), 43 (1996)

A53) Hummel, K., Sputtek, A., Kühnl, P.

Influence of MNC concentration during cryopreservation on progenitor cell recovery

Infusionsther. Transfusionsmed. 23 (Suppl 3), 42 (1996)

A52) Sputtek, A., Roos, D., Lubitz, B., Gutensohn, K., Rapp, Ch., Kühnl, P.

Einfluß des Bestrahlungsvorgangs auf gefilterte Erythrozyten-Konzentrate in additiver  
Lösung (PAGGS-M)

Infusionsther. Transfusionsmed. 23 (Suppl 3), 15 (1996)

A51) Sputtek, A., Horn, E.-P., Voll, M., Hummel, K., Hiller, J., Lüdemann, K., Schulte  
am Esch, J., Kühnl, P.

Erste Ergebnisse der Klinischen Phase-II-Studie zur Sicherheit und Verträglichkeit  
autologer, mittels Hydroxyethylstärke (HES) kryokonservierter Erythrozyten (RBC)

Infusionsther. Transfusionsmed. 23 (Suppl 3), 5 (1996)

A50) Gutensohn, K., Naujokat C., Sputtek, A., Cassens, U., Roos, D., Kühnl, P.

Decrease in absolute and relative counts of natural killer cells (CD3-/CD16+CD56+)  
during preoperative autologous blood donation

Infusionsther. Transfusionsmed. 23 (Suppl 3) 2 (1996)

A49) Sputtek, A., Jetter, S., Hummel, K., Kühnl, P.

How much DMSO and which freezer do we need? - Influence of cooling rate and  
composition of the cryoprotective solution on the recovery of frozen/thawed  
peripheral blood stem cells

Exp. Hematol. 24, 1027 (1996)

A48) Sputtek, A., Jetter, S., Hummel, K., Löliger, C., Kühnl, P.

Interaction of HES/DMSO concentration and cooling rate on the recovery of cryopreserved human PBSC  
Cryobiology 33, 650-651 (1996)

A47) Sputtek, A., Horn, E.-P., Voll, M., Hummel, K., Hiller, J., Lüdemann, K., Schulte am Esch, J., Kühnl, P.

A clinical study to determine the safety and tolerance of HES-cryopreserved autologous red blood cells  
Cryobiology 33, 650-651 (1996)

A46) Horn, E.-P., Sputtek, A., Rudolf, B., Standl, Th.

Sicherheit und Toleranz bei autologer Transfusion von HES-kryokonservierten Erythrozyten-Konzentraten  
Anaesthesist 45 (Suppl 2) 130 (1996)

A45) Wittkopf, D., Sputtek, A., Feddern, P., Reckhaus, M.-L., Kühnl, P.  
Vergleich von 4 Methoden zur Kryokonservierung von Testerythrozyten  
Infusionsther. Transfusionsmed. 22 (Suppl 2) 10 (1995)

A44) Langer, R., Herold, T., Sputtek, A., Henrich, H.A.

Lipidperoxidation als eine Hämolyseursache bei mit Hydroxyethylstärke (HES) kryokonservierten Erythrozyten  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 8 (1995)

A43) Hummel, K., Zeller, W., Gutensohn, K., Sputtek, A., Stockschläder, M., Zander, A., Kühnl, P.

PBSC Mobilisation und -Separation bei gesunden Spendern  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 43 (1995)

A42) Gutensohn, K., Sputtek, A., Cassens, U., Caspers, U., Kühnl, P.

Optimized storage conditions for flow cytometric analysis of peripheral blood stem cells  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 58 (1995)

A41) Voll, M., Sputtek, A., Wittkopf, D., Kühnl, P.

Kryokonservierung von McLeod-Erythrozyten mit Hydroxyethylstärke (HES)  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 11 (1995)

A40) Sputtek, A., Gutensohn, K., Langer, R., Hammes, R., Warnken, U.H., Henrich, H.A., Kühnl, P.

Elimination von Hydroxyethylstärke (HES) nach autologer Transfusion von damit kryokonservierten Erythrozyten  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 10 (1995)

A39) Sputtek, A., Lubitz, B., Naujokat, C., Zinke, H., Roos, D., Kühnl, P.

Welchen Einfluß hat die Unterbrechung der Kühlkette bei der Lagerung von Erythrozyten-Konzentraten nach dem "Eppendorfer-Modell"?  
Infusionsther. Transfusionsmed. 22 (Suppl 2), 10 (1995)

A38) Sputtek, A., Meuser, D., Kühnl, P.

A method for freezing red blood cells for transfusion using hydroxyethyl starch (HES)

Cryobiology 32, 544-545 (1995)

A37) Gutensohn, K., Bartsch, N., Cassens, U., Sputtek, A., Ganschow, I., Kühnl, P.  
Platelet activation (PA) profiles in donors following cytappheresis: A flow cytometric follow-up for 4 days  
Ann. Hematol. 70 (Suppl 1), 11 (1995)

A36) Gutensohn, K., Bartsch, N., Sputtek, A., Ganschow, I., Cassens, U., Kühnl, P.  
Flow cytometry as a tool to determine cytappheresis-induced platelet activation (PA)  
Ann. Hematol. 70 (Suppl 1), 10 (1995)

A35) Sputtek, A.  
Cryopreservation of human red blood cells: A comparison of the cryoprotectants gelatine, dextran, and hydroxyethyl starch  
Cryo Letters 16, 58 (1995)

A34) Spieles, G., Kresin, M., Loges, K., Sputtek, A., Heschel, I., Rau, G.  
Hemolysis and devitrification in erythrocytes cryopreserved with hydroxyethyl starch  
Cryo Letters 16, 58 (1995)

A33) Meuser, D., Hoevel, M., Sputtek, A.  
Full unit red blood cell (RBC) freezing in the presence of 12% hydroxyethyl starch (HES)  
Cryo Letters 16, 58 (1995)

A32) Sputtek, A., Sensendorf, H., Fründ, V., Birmanns, H., Kühnl, P.  
Transfusion of 3 units of autologous RBC frozen in the presence of HES without postthaw washing - a case report  
Infusionsther. Transfusionsmed. 21 (Suppl 2), 74 (1994)

A31) Sputtek, A.  
An optimized procedure for the cryopreservation of human red blood cells (RBC) using hydroxyethyl starch (HES) and liquid nitrogen (LN2)  
Infusionsther. Transfusionsmed. 21 (Suppl 2), 74 (1994)

A30) Sputtek, A., Barthel, B., Kühnl, P.  
A 4% HES method for the cryopreservation of human platelets compared with the standard 5% DMSO method  
Infusionsther. Transfusionsmed. 21 (Suppl 2), 74 (1994)

A29) Langer, R., Albrecht, R., Hempel, K., Krug, S., Sputtek, A., Steigerwald, R., Trenkel, K., Henrich, H.A.  
Studies on 24-hour post-transfusion survival and long-term survival of autologous with hydroxyethyl starch (HES) cryopreserved erythrocytes in dog  
Infusionsther. Transfusionsmed. 21 (Suppl 2), 72-73 (1994)

A28) Sputtek, A., Heschel, I., Kühnl, P.  
Tiefemperaturkonservierung menschlicher roter Blutkörperchen mit dem Kryoprotektiv Hydroxyethylstärke (HES) - Bedeutung der Abkühlrate  
Chem. Ing. Tech. 66, 1217-1218 (1994)

- A27) Sputtek, A., Sensendorf, H., Freund, V., Birmanns, H., Kühnl, P.  
Successful transfusion of previously frozen autologous red blood cells without postthaw washing  
Cryobiology 31, 584-585 (1994)
- A26) Sputtek, A., Warnken, U.H., Langer, R., Henrich, H.A., Kühnl, P.  
Hydroxyethyl starch plasma concentration profiles after administration of HES protected frozen erythrocytes  
Cryobiology 31, 584 (1994)
- A25) Rowe, A.W., Sputtek, A.  
Cryopreservation of red blood cells for transfusion  
Cryobiology 31, 583 (1994)
- A24) Sputtek, A., Barthel, B., Kühnl, P.  
Comparison of a 4% hydroxyethyl starch (HES) method with the 5% dimethyl sulfoxide technique for the cryopreservation of platelet concentrates (PC)  
Vox Sang. 67 (Suppl 2), 33 (1994)
- A23) Sputtek, A., Singbartl, G., Schleinzner, W., Kühnl, P.,  
Cryopreservation of human red blood cells (RBC) using hydroxyethyl starch (HES) and liquid nitrogen: Autologous retransfusion in healthy volunteers  
Vox Sang. 67 (Suppl 2), 32 (1994)
- A22) Sputtek, A., Sensendorf, H., Fründ, V., Birmanns, H., Kühnl, P.  
Substitution von Blutvolumen und Sauerstofftransportkapazität durch Autologe, mit Hydroxyethylstärke (HES) kryokonservierte Erythrozyten im Ein-Schritt-Verfahren - Ein Fallbericht  
Anaesthesist 43 (Suppl 1), 141 (1994)
- A21) Sputtek, A., Roling, Ch., Singbartl, G., Schleinzner, W., Rau, G.  
First autologous retransfusion of human red blood cells (450 ml) cryopreserved in the presence of 11.5% HES without postthaw washing  
Cryobiology 30, 657 (1993)
- A20) Sputtek, A., Barthel, B., Rau, G.  
Cryopreservation of human platelets: An in vitro comparison of a 4% HES method with the standard 5% DMSO method  
Cryobiology 30, 656-657 (1993)
- A19) Sputtek, A., Heschel, I., Bryant, G.  
Cryoprotection of blood cells by polymeric compounds: A discussion of possible mechanisms  
Cryobiology 30, 654-655 (1993)
- A18) Sputtek, A., Gier, M.  
Osmotic properties of highly concentrated ternary HES-NaCl-H<sub>2</sub>O solutions  
Cryobiology 30, 621 (1993)
- A17) Sputtek, A., Singbartl, G., Schleinzner, W., Rau, G.

Kryokonservierung von Erythrozyten mit Hydroxyethylstärke (HES): In-vivo-Ergebnisse im autologen Retransfusionsmodell beim Menschen  
Infusionsther. 20 (Suppl 3), 62 (1993)

A16) Sputtek, A., Hammes, R., Warnken, U.H., Langer, R., Henrich, H.A., Rau, G.  
Konzentrationsverlauf von Hydroxyethylstärke (HES) in Blut und Urin nach autologer Retransfusion von damit kryokonservierten Erythrozyten mittels Hochleistungs-Gelpermeationschromatographie  
Anaesthesist 42 (Suppl 1), 406 (1993)

A15) Sputtek, A., Singbartl, G., Schleinzer, W., Rau, G.  
Welche Faktoren begrenzen die Transfusion von mittels Hydroxyethylstärke (HES) kryokonservierten autologen Erythrozytenkonzentraten?  
Anaesthesist 42 (Suppl 1), 276 (1993)

A14) Sputtek, A., Singbartl, G., Schleinzer, W., Rau, G.  
Autologous transfusion of human erythrocytes cryopreserved with hydroxyethyl starch (HES) in seven volunteers  
Cryobiology 29, 766-767 (1992)

A13) Sputtek, A.  
Cryopreservation of various types of blood cells: What can we learn from the past 10 years?  
Cryobiology 29, 765 (1992)

A12) Sputtek, A., Warnken, U.H.  
The effect of hydroxyethyl starch (HES) molecular weight on the amount of uncrystallized water  
Cryobiology 29, 707-708 (1992)

A11) Sputtek, A., Singbartl, G., Schleinzer, W., Rau, G.  
Autologe Retransfusion von hydroxyethylstärkehaltigen, tiefgekühlten Erythrozyten-Konzentraten im Selbstversuch  
Anaesthesist 41 (Suppl 1), 162 (1992)

A10) Langer, R., Kron, W., Schmid, H., Sputtek, A., Steigerwald, R., Trenkel, K., Rau, G., Henrich, H.A.  
Studies on functional acceptability of autologous with hydroxyethyl starch (HES) cryopreserved erythrocytes in dogs  
Clin. Hemorheol. 12, 476 (1992)

A9) Sputtek, A., Dreyer, M., Heschel, I., Rau, G.  
Survival of human red blood cells (RBC) after cryopreservation as affected by the molecular weight of the cryoprotectant hydroxyethyl starch (HES)  
Cryobiology 28, 554 (1991)

A8) Sputtek, A., Schmid, H., Henrich, H.A., Körber, Ch., Rau, G.  
Cryopreservation of dog red blood cells (DRBC) with hydroxyethyl starch (HES) for autologous retransfusion as an animal study  
Cryobiology 28, 546 (1991)

A7) Sputtek, A., Körber, Ch., Langer, R., Schmid, H., Steigerwald, R., Trenkel, K., Kron, W., Henrich, H.A., Rau, G.  
Kryokonservierung von Erythrozyten mit Hydroxyethylstärke (HES): In-vitro-Ergebnisse auf dem Weg zum autologen Retransfusionsmodell beim Hund  
Infusionsther. 18 (Suppl 2), 37 (1991)

A6) Sputtek, A., Körber, Ch., Rau, G.  
Cryopreservation of human erythrocytes with hydroxyethylated starches under variation of starch modification and concentration, electrolyte content, hematocrit, and cooling rate  
Cryobiology 27, 667-668 (1990)

A5) Sputtek, A.  
A more detailed understanding of molecular properties of etherified starches used as cryoprotectants  
Cryobiology 27, 632 (1990)

A4) Sputtek, A., Körber, Ch., Mayr, W.R., Rau, G.  
Cryopreservation of human platelets: A comparison of different viability assays  
Cryobiology 25, 524 (1988)

A3) Sputtek, A., Körber, Ch., Bacher, Ch., Rau, G.  
Further improvement of the cryopreservation of human red blood cells with hydroxyethyl starch  
Cryobiology 25, 523 (1988)

A2) Manseck, A., Sputtek, A., Körber, Ch.  
The influence of the cryoprotectants trehalose and dimethyl sulfoxide on granulocyte function  
Cryobiology 25, 515-516 (1988)

A1) Körber, Ch., Hartmann, U., Brohm, A., Classen, I., Scheiwe, M.W., Sputtek, A., Rau, G.  
Cryopreservation of human platelets with hydroxyethyl starch  
Cryobiology 23, 576 (1986)